



AgroSym
2019

BOOK OF

PROCEEDINGS



*X International Scientific Agriculture Symposium
"AGROSYM 2019"
Jahorina, October 03-06, 2019*

BOOK OF PROCEEDINGS

**X International Scientific Agriculture Symposium
“AGROSYM 2019”**



Jahorina, October 03 - 06, 2019

Impressum

X International Scientific Agriculture Symposium „AGROSYM 2019“

Book of Abstracts Published by

University of East Sarajevo, Faculty of Agriculture, Republic of Srpska, Bosnia
University of Belgrade, Faculty of Agriculture, Serbia
Mediterranean Agronomic Institute of Bari (CIHEAM - IAMB) Italy
International Society of Environment and Rural Development, Japan
Balkan Environmental Association (B.EN.A), Greece
Centre for Development Research, University of Natural Resources and Life Sciences (BOKU), Austria
Perm State Agro-Technological University, Russia
Voronezh State Agricultural University named after Peter The Great, Russia
Faculty of Bioeconomy Development, Vytautas Magnus University, Lithuania
Selçuk University, Turkey
University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
Slovak University of Agriculture in Nitra, Slovakia
Ukrainian Institute for Plant Variety Examination, Kyiv, Ukraine
National University of Life and Environmental Sciences of Ukraine, Kyiv, Ukraine
Valahia University of Targoviste, Romania
National Scientific Center „Institute of Agriculture of NAAS“, Kyiv, Ukraine
Saint Petersburg State Forest Technical University, Russia
University of Valencia, Spain
Faculty of Agriculture, Cairo University, Egypt
Tarbiat Modares University, Iran
Chapingo Autonomous University, Mexico
Department of Agricultural, Food and Environmental Sciences, University of Perugia, Italy
Higher Institute of Agronomy, Chott Mariem-Sousse, Tunisia
Watershed Management Society of Iran
Institute of Animal Science- Kostinbrod, Bulgaria
Faculty of Agriculture, University of Banja Luka, Bosnia and Herzegovina
Faculty of Economics Brcko, University of East Sarajevo, Bosnia and Herzegovina
Biotechnical Faculty, University of Montenegro, Montenegro
Institute of Field and Vegetable Crops, Serbia
Institute of Lowland Forestry and Environment, Serbia
Institute for Science Application in Agriculture, Serbia
Agricultural Institute of Republic of Srpska - Banja Luka, Bosnia and Herzegovina
Maize Research Institute “Zemun Polje”, Serbia
Faculty of Agriculture, University of Novi Sad, Serbia

Editor in Chief

Dusan Kovacevic

Technical editors

Sinisa Berjan

Milan Jugovic

Noureddin Driouech

Rosanna Quagliariello

Website:

<http://agrosym.ues.rs.ba>

CIP - Каталогизacija u publikaciji
Народна и универзитетска библиотека
Републике Српске, Бања Лука

631(082)

INTERNATIONAL Scientific Agricultural Symposium "Agrosym 2019" (10)
(Jahorina)

Book of Proceedings [Elektronski izvor] / X International Scientific Agriculture
Symposium "Agrosym 2019", Jahorina, October 03 - 06, 2019 ; [editor in chief Dušan
Kovačević]. - East Sarajevo : Faculty of Agriculture, 2019

Način pristupa (URL): <http://agrosym.ues.rs.ba/index.php/en/archive>. -
Библиографија уз радове. - Регистар.

ISBN 978-99976-787-2-0

COBISS.RS-ID 8490776

X International Scientific Agricultural Symposium “Agrosym 2019”
Jahorina, October 03-06, 2019, Bosnia and Herzegovina

HONORARY COMMITTEE

BORIS PASALIC, Minister of Agriculture, Water Management and Forestry of Republic of Srpska, Bosnia, SRDJAN RAJCEVIC, Minister of Scientific-Technological Development, Higher Education and Information Society of Republic of Srpska, Bosnia, MILAN KULIC, Rector of the University of East Sarajevo, Bosnia, DUSAN ZIVKOVIC, Dean of the Faculty of Agriculture, University of Belgrade, Serbia, MAURIZIO RAELI, Director of the Mediterranean Agronomic Institute of Bari, Italy, MARIO T. TABUCANON, President of the International Society of Environment and Rural Development, Japan, FOKIAON K. VOSNIAKOS, President of the Balkan Environmental Association (B.EN.A), Greece, MUSTAFA ŞAHIN, Rector of the Selcuk University, Turkey, ALEKSEY ANDREEV, Rector of the Perm State Agro-Technological University, Russia, NIKOLAY I. BUKHTOYAROV, Rector of the Voronezh State Agricultural University named after Peter The Great, Russia, ANTANAS MAZILIAUSKAS, Rector of the Vytautas Magnus University Agriculture Academy, Lithuania, BARBARA HINTERSTOISSER, Vice-Rector of the University of Natural Resources and Life Sciences (BOKU), Austria, JOSÉ SERGIO BARRALES DOMÍNGUEZ, Rector of the Chapingo Autonomous University, Mexico, KLAUDIA HALÁSZOVÁ, Rector of the Slovak University of Agriculture in Nitra, Slovakia, CALIN D. OROS, Rector of the Valahia University of Targoviste, Romania, VIKTOR KAMINSKYI, Director of National Scientific Center „Institute of Agriculture of NAAS“, Kyiv, Ukraine, AMR AHMED MOSTAFA, Dean of the Faculty of Agriculture, Cairo University, Egypt, SORIN MIHAI CIMPEANU, Rector of the University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania, FRANCESCO TEI, Director of the Department of Agricultural, Food and Environmental Sciences, University of Perugia, Italy, DAVUT KARAYEL, Dean of Faculty of Agriculture, University of Akdeniz - Antalya, Turkey, CHOKRI THABET, the General Director of the High Agronomic Institute of Chott Mariem, Sousse, Tunisia, SEYED HAMIDREZA SADEGHI, Professor at Tarbiat Modares University and the President of the Watershed Management Society of Iran, Iran, IVAN YANCHEV, Director of the Institute of Animal Science- Kostinbrod, Bulgaria, ZLATAN KOVACEVIC, Dean of the Faculty of Agriculture, University of Banja Luka, Bosnia and Herzegovina, LAZAR RADOVANOVIC, Dean of the Faculty of Economics Brcko, Universtiy of East Sarajevo, Bosnia and Herzegovina, BOZIDARKA MARKOVIC, Dean of the Biotechnical Faculty, University of Podgorica, Montenegro, SNEZANA JANKOVIC, Director of the Institute for Science Application in Agriculture, Serbia, SASA ORLOVIC, Director of the Institute of Lowland Forestry and Environment, Serbia, BRANKO KOVACEVIC, President of the Academy of Engineering Sciences of Serbia, Serbia, VOJISLAV TRKULJA, Director of Agricultural Institute of Republic of Srpska - Banja Luka, Bosnia and Herzegovina, BRANKA KRESOVIC, Director of the Maize Research Institute “Zemun Polje”, Serbia, SVETLANA BALESEVIC-TUBIC, Director of the Institute of Field and Vegetable Crops, Serbia, NEDELJKO TICA, Dean of the Faculty of Agriculture, University of Novi Sad, Serbia, RODNE NASTOVA, Director of the Institute for Animal Science, Skoplje, Macedonia, RADOVAN PEJANOVIC, President of Balkan Scientific Association of Agricultural Economics, Serbia, JONEL SUBIC, Director of the Institute of Agricultural Economics, Serbia

SCIENTIFIC COMMITTEE

DUSAN KOVACEVIC, Faculty of Agriculture, University of Belgrade, Serbia, Prof. dr William Meyers, Howard Cowden Professor of Agricultural and Applied Economics, University of Missouri, USA, JOHN BRAYDEN, Norwegian Agricultural Economics Research Institute (NILF), Norway, STEVE QUARIE, Visiting Professor, School of Biology, Newcastle University, United Kingdom, ANDREAS MELCHER, CDR, University of Natural Resources and Life Sciences (BOKU), Vienna, Austria, DANI SHTIENBERG, full professor, Department of Plant pathology and Weed Research, ARO, the Volcani Center, Bet Dagan, Israel, THOMAS G. JOHNSON, University of Missouri – Columbia, USA, DIETER TRAUTZ, University of Applied Science, Germany, MACHITO MIHARA, Tokyo University of Agriculture, Japan, MARKUS SCHERMER, Department of Sociology, University of Innsbruck, Austria, JORGE BATLLE-SALES, Department of Biology, University of Valencia, Spain, SERGEI ELISEEV, Vice-Rector for Research and Innovations, Perm State Agro-Technological University, Russia, RICHARD BARICHELLO, Faculty of Land and Food Systems, University of British Columbia, Canada, NOVO PRZULJ, Faculty of Agriculture, University of Banjaluka, Bosnia and Herzegovina, TATIANA SIVKOVA, Faculty for Veterinarian Medicine and Zootechny, Perm State Agro-Technological University, Russia, ADRIANO CIANI, Department of Agricultural, Foods and Environmental Sciences, Perugia University, Italy, ALEKSEJ LUKIN, Voronezh State Agricultural University named after Peter The Great, Russia, MATTEO VITTUARI, Faculty of Agriculture, University of Bologna, Italy, SEYED MOHSEN HOSSEINI, Faculty of Natural Resources, Tarbiat Modares University, Iran, ARDIAN MACI, Faculty of Agriculture and Environment, Agricultural University of Tirana, Albania, REGUCIVILLA A. POBAR, Bohol Island State University, Philippines, SUDHEER KUNDUKULANGARA PULISSERY, Kerala Agricultural University, India, EPN UDAYAKUMARA, Faculty of Applied Sciences, Sabaragamuwa University, Sri Lanka, VLADIMIR SMUTNÝ, full professor, Mendel University, Faculty of agronomy, Czech Republic, FRANC BAVEC, full professor, Faculty of Agriculture and Life Sciences, Maribor, Slovenia, JAN MOUDRY, full professor, Faculty of Agriculture, South Bohemia University, Czech Republic, STEFAN TYR, full professor, Faculty of Agro-biology and Food Resources, Slovakia, NATALIJA BOGDANOV, Faculty of Agriculture, University of Belgrade, Serbia, SABAHUDIN BAJRAMOVIC, Faculty of Agriculture and Food Sciences, University of Sarajevo, Bosnia, FRANCESCO PORCELLI, University of Bari Aldo Moro, Italy, VASILJE ISAJEV, Faculty of Forestry, University of Belgrade, Serbia, ELAZAR FALLIK, Agricultural Research Organization (ARO), Volcani, Israel, JUNAID ALAM MEMON, Pakistan Institute of Development Economics, Pakistan, HIROMU OKAZAWA, Faculty of Regional Environment Science, Tokyo University of Agriculture, Japan, PANDI ZDRULI, Land and Water Resources Department; IAMB, Italy, MLADEN TODOROVIC, Land and Water Resources Department; IAMB, Italy, HAMID EL BILALI, CDR, University of Natural Resources and Life Sciences (BOKU), Vienna, Austria, MAKSYM MELNYCHUK, National Academy of Agricultural Science of Ukraine, Ukraine, BORYS SOROCHYNSKYI, Ukrainian Institute for Plant Variety Examination, Kyiv, Ukraine, LORENZ PROBST, CDR, University of Natural Resources and Life Sciences (BOKU), Vienna, Austria, MOHSEN BOUBAKER, High Institute of Agronomy of Chott

Meriem, Sousse, Tunisia, NOUREDDIN DRIOUECH, Coordinator of MAIB Alumni Network (FTN), Mediterranean Agronomic Institute of Bari, Italy, ION VIOREL, University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania, CHULEEMAS BOONTHAI IWAI, Faculty of Agriculture, Khon Kaen University, Thailand, WATHUGE T.P.S.K. SENARATH, Department of Botany, University of Sri Jayewardenepura, Colombo, Sri Lanka, HAMADA ABDELRAHMAN, Soil Science Dept., Faculty of Agriculture, Cairo University, Egypt, MAYA IGNATOVA, Agricultural Academy – Sofia, Bulgaria, IOANNIS N. XYNIAS, School of Agricultural Technology & Food Technology and Nutrition, Western Macedonia University of Applied Sciences, Greece, ARTUR RUTKIEWICZ, Department of Forest Protection, Forest Research Institute - IBL, Poland, MOHAMMAD SADEGH ALLAHYARI, Islamic Azad University, Rasht Branch, Iran, LALITA SIRIWATTANANON, Faculty of Agricultural Technology, Rajamangala University of Technology Thanyaburi (RMUTT), Thailand, KONSTANTIN KORLYAKOV, Perm Agricultural Research Institute, Russia, MOHAMMAD FAROOQUE HASSAN, Shaheed Benazir Bhutto University of Veterinary & Animal Sciences Sakrand, Sindh, Pakistan, LARYSA PRYSIAZHNIUK, Ukrainian Institute for Plant Variety Examination, Kyiv, Ukraine, OKSANA KLIACHENKO, National University of Life and Environmental Science of Ukraine, Ukraine, IVAN SIMUNIC, Department of amelioration, Faculty of agriculture, University of Zagreb, Croatia, ABID HUSSAIN, International Centre for Integrated Mountain Development (ICIMOD), Nepal, AMRITA GHATAK, Gujarat Institute of Development Research (GIDR), India, NASER SABAGHNI, University of Maragheh, Iran, MONICA PAULA MARIN, Department for Animal Husbandry, University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania, PENKA MONEVA, Institute of Animal Science - Kostinbrod, Bulgaria, MOSTAFA K. NASSAR, Animal husbandry Dept., Faculty of Agriculture, Cairo University, Egypt, MÁRTA BIRKÁS, full professor, St. Istvan University, Godollo – Hungary, ANDRZEJ KOWALSKI, Director of the Institute for Agricultural and Food Economy, Warszawa-Poland, YALCIN KAYA, The Director of the Plant Breeding Research Center, University of Trakya, Turkey, SANJA RADONJIC, Biotechnical Faculty, University of Montenegro, Montenegro, IONELA DOBRIN, Department for Plant Protection, University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania, INOCENCIO BUOT JR., Institute of Biological Sciences, College of Arts and Sciences, University of the Philippines Los Banos, Philippines, KAROL WAJSZCZUK, Poznan University of Life Sciences, Poland, REDOUANE CHOUKR-ALLAH, International Center for Biosaline Agriculture (ICBA), United Arab Emirates, MOHAMMAD AL-MAMUN, Department of Animal Nutrition, Bangladesh Agricultural University, Bangladesh, ANUCHA WITTAYAKORN-PURIPUNPINYOO, School of Agriculture and Co-operatives, Sukhothai Thammathirat Open University, Nonthaburi, Thailand, NEDELJKA NIKOLOVA, Institute for Animal Science, Ss. Cyril and Methodius University in Skopje, Republic of Macedonia, IGNACIO J. DÍAZ-MAROTO, High School Polytechnic, University of Santiago de Compostela, Spain, NIDAL SHABAN, University of Forestry Sofia, Bulgaria, YOUSSEF SASSINE, Lebanese University Beirut, Lebanon, CAFER TOPALOGLU, Faculty of Tourism, Mugla Sitki Kocman University, Turkey, SEYED HAMIDREZA SADEGHI, Faculty of Natural Resources, Tarbiat Modares University, Iran, MOHSEN MOHSENI SARAVI, University of Teheran and Member of WMSI Management Board, Iran, MAHMOOD ARABKHEDRI, Soil Conservation and Watershed Management Research Institute and Member of WMSI Management Board, Iran, ATAOLLAH KAVIAN, Sari Agricultural Science and Natural Resources University and Member of WMSI Management Board, Iran, TUGAY AYASAN, Department of Organic Farming Business Management, Osmaniye, Applied Science School of Kadirli, Osmaniye Korkut Ata University, Turkey, SAKINE ÖZPINAR, Department of Farm Machinery and Technologies Engineering, Faculty of Agriculture, Çanakkale Onsekiz Mart University, Çanakkale, Turkey, SHEREIN SAEIDE ABDELGAYED, Faculty of Veterinary Medicine, Cairo University, Cairo, Egypt, ZOHREH MASHAK, Islamic Azad University, Karaj Branch, Iran, KHALID AZIM, National Institute of Agriculture Research, Morocco, SRDJAN LALIC, Vice-dean University of East Sarajevo, Bosnia and Herzegovina, ZELJKO VASKO, Faculty of Agriculture, University of Banja Luka, Bosnia and Herzegovina, KUBILAY BAŞTAŞ, Department of Plant Protection, Faculty of Agriculture, Selçuk University, Turkey, BRANKA KRESOVIC, Director of the Maize Research Institute “Zemun Polje”, Serbia, NENAD DELIC, Maize Research Institute “Zemun Polje”, Serbia, MILAN STEVANOVIC, Maize Research Institute “Zemun Polje”, Serbia, SVETLANA BALESEVIC-TUBIC, Institute of Field and Vegetable Crops Novi Sad, Serbia, ANA MARJANOVIC JEROMELA, Institute of Field and Vegetable Crops Novi Sad, Serbia, TATJANA ZDRALIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, ALEKSANDRA GOVEDARICA-LUCIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, DESIMIR KNEZEVIC, University of Pristina, Faculty of Agriculture, Kosovska Mitrovica - Lesak, Kosovo i Metohija, Serbia, SNEZANA MLADENOVIC-DRINIC, Maize Research Institute “Zemun Polje”, Serbia, NEBOJSA MOMIROVIC, Faculty of Agriculture, University of Belgrade, Serbia, VELIBOR SPALEVIC, Faculty of Philosophy, Geography, University of Montenegro, ZORAN JOVOVIC, Biotechnical Faculty, University of Montenegro, Montenegro, DANIJEL JUG, associate professor, Faculty of Agriculture, University of Osijek, Croatia, MILAN MARKOVIC, Biotechnical Faculty, University of Montenegro, Montenegro, ZELJKO DOLIJANOVIC, Faculty of Agriculture, University of Belgrade, Serbia, DEJAN STOJANOVIC, Institute of Lowland Forestry and Environment, Serbia, DOBRIVOJ POŠTIĆ, Institute for plant protection and environment, Belgrade, Serbia, SRDJAN STOJNIC, Institute of Lowland Forestry and Environment, Serbia

ORGANIZING COMMITTEE

VESNA MILIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, DEJAN BOKONJIC, Vice rector of the University of East Sarajevo, Bosnia, DEJANA STANIC, Dean of the Faculty of Agriculture, University of East Sarajevo, Bosnia, MAROUN EL MOUJABBER, Mediterranean Agronomic Institute of Bari, Italy, ROSANNA QUAGLIARIELLO, Mediterranean Agronomic Institute of Bari, Italy, NOUREDDIN DRIOUECH, Coordinator of MAIB Alumni Network (FTN), Mediterranean Agronomic Institute of Bari, Italy, ALEKSANDRA DESPOTOVIC, Biotechnical Faculty Podgorica, University of Montenegro, Montenegro, MILIC CUROVIC, The journal “Agriculture and Forestry”, Biotechnical Faculty Podgorica, University of Montenegro, Montenegro, ANA MARJANOVIC JEROMELA, Institute of Field and Vegetable Crops, Serbia, OKSANA FOTINA, International Relations Center, Perm State Agro-Technological University, Russia, TATIANA LYSK, International Relations Office, Voronezh State Agricultural University named after Peter The Great, Russia, ANASTASIJA NOVIKOVA, Faculty of Bioeconomy Development, Vytautas Magnus University, Lithuania, ENGR. TEODORA POPOVA, Institute of Animal Science - Kostinbrod, Bulgaria, MEHMET MUSA OZCAN, Faculty of Agriculture, Selçuk University, Turkey, LAZAR

RADOVANOVIC, Faculty of Economics Brcko, University of East Sarajevo, Bosnia and Herzegovina, NIKOLA PACINOVSKI, Institute for Animal Science, Ss. Cyril and Methodius University in Skopje, Republic of Macedonia, ABDULVAHED KHALEDI DARVISHAN, Faculty of Natural Resources, Tarbiat Modares University, Iran, HAMADA ABDELRAHMAN, Soil Science Dept., Faculty of Agriculture, Cairo University, Egypt, ECATERINA STEFAN, University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania, JEERANUCH SAKKHAMDUANG, The International Society of Environmental and Rural Development, Japan, RAOUDHA KHANFIR BEN JENANA, High Institute of Agronomy of Chott Meriem, Sousse, Tunisia, ERASMO VELÁZQUEZ CIGARROA, Department of Rural Sociology, Chapingo Autonomous University, Mexico, VEDRAN TOMIC, Institute for Science Application in Agriculture, Serbia, MILAN STEVANOVIC, Maize Research Institute "Zemun Polje", Serbia, ANDREJ PILIPOVIC, Institute of Lowland Forestry and Environment, Serbia, NIKOLA PUVACA, Faculty of Economics and Engineering Management University of Business Academy, Serbia, MORTEZA BEHZADFAR, Tarbiat Modares University, Tehran, Iran, LARYSA PRYSIAZHNIUK, Ukrainian Institute for Plant Variety Examination, Kyiv, Ukraine, BRANISLAVKA BOROJA, Agricultural Institute of Republic of Srpska - Banja Luka, Bosnia, BILJANA GRUJIC, Institute of Agriculture Economics, Serbia, MARKO GUTALJ, Faculty of Agriculture, University of East Sarajevo, Bosnia, MIROSLAV LALOVIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, MILAN JUGOVIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, IGOR DJURDJIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, MILENA STANKOVIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, STEFAN STJEPANOVIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, STEFAN BOJIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, TANJA JAKISIC, Faculty of Agriculture, University of East Sarajevo, Bosnia, TIJANA BANJANIN, Faculty of Agriculture, University of East Sarajevo, Bosnia, SINISA BERJAN, Faculty of Agriculture, University of East Sarajevo, Bosnia, General secretary

EFFECT OF DRYING ON THE CHANGE OF SUGAR CONTENT IN PLUM FRUITS

Olga MITROVIĆ^{1*}, Branko POPOVIĆ¹, Nemanja MILETIĆ², Aleksandar LEPOSAVIĆ¹,
Aleksandra KORICANAC¹

¹Fruit Research Institute, Čačak, Serbia

²University of Kragujevac, Faculty of Agriculture in Čačak, Serbia

*Corresponding author: omitrovic@institut-cacak.org

Abstract

Drying is one of the oldest methods of preserving fruit. In the course of drying process, there is a change in some components of chemical composition to a degree depending on food type, its composition, and drying process itself. The testing was performed in three replications using plum fruits of the 'Čačanska Lepotica', 'Mildora', 'Čačanska Rodna' and 'Stanley' cultivars, at the optimum ripening stage required for drying, based on soluble solid contents. Drying was performed at the experimental dryer using the convective (streaming) drying process, at two constant air temperatures, 90 °C and 70 °C, until attaining 75% of total dry matter in the prunes. In addition to the control, the fruits were subjected to a pretreatment consisting of dipping in boiling water. This paper reviews the values of total sugars, invert sugars and sucrose in fruits of fresh and dried plums (calculated in grams per 100 grams of total dry matter), as well as the change of these parameters in prunes in relation to the fresh fruits (expressed in percentages). Drying temperature affected the change of total sugars, invert sugars and sucrose content in fruit of the tested plum cultivars. On the other side, dipping as an applied pretreatment had no effect on the change of these parameters except in the cultivar 'Čačanska Rodna', which dipped fruits are found to have higher decrease of sucrose compared with non-treated fruits. During drying, hydrolysis of sucrose occurred, manifesting in a dramatic decrease of its content in dried fruits, in relation to the starting raw material in all of the tested cultivars. Intensity of changes was conditioned by varietal characteristics.

Key words: *prune, drying temperature, dipping, invert sugar, sucrose.*

Introduction

Plum (*Prunus domestica* L.) is considered a national fruit species ranking first in the fruit growing of Serbia by both number of trees and by its annual production. Although a major part of plum yield in Serbia is processed into brandy (Popović et al., 2006), the most significant product coming from plum, in terms of nutritional value, is prune. Prune represents significant energy source for human body though is more commonly classified in high nutritional food with special dietetic and physiological importance, primarily due to its protective and therapeutic effect on digestive tract of consumers (Zlatković, 2000; Piga et al., 2003).

Drying is defined as a process of moisture loss resulting from simultaneous existence of the heat and mass transfer phenomena. Plum drying is a slow and long-term process because whole fruits are dried, hence, there are real conditions for various chemical transformations, primarily carbons, since they make up over 90% of the total dry matter of fruit (Friedman, 1996). In plum fruit, the most predominant are sugars, i.e. fermentable sugars (fructose, glucose and sucrose). During plum drying, in addition to oxidation reaction and acid hydrolysis (Dikerman et al., 2004), non-enzymatic browning reactions with sugars participating, such as caramelization and Maillard reaction (Manzocco et al., 2001; Kim end Lee, 2008), can be conducted too. Depending on the intensity of these reactions, characteristic colour and flavour of prune appears but on the other hand, they may lead to degradation of

product aroma and appearance (Dikerman et al., 2004; Sanz et al., 2001). Appropriate drying technology and raw material can affect the flow of these reactions in order to minimize the change of product quality, which is an ultimate goal of each processing.

The aim of the work was to examine the effect of drying temperature and dipping on the change of sugar content, both total and invert sugars as well as sucrose content in fruits of the plum cultivars 'Čačanska Lepotica', 'Mildora', 'Čačanska Rodna' and 'Stanley'.

Material and Methods

For the purpose of testing, plum fruits of the cultivars 'Čačanska Lepotica', 'Mildora', 'Čačanska Rodna' and 'Stanley' were obtained from the plantations of the Fruit Research Institute Čačak, in which standard agro- and pomotechnical practices were performed. Fruits for drying were selectively picked at full maturity stage for the appropriate cultivar, with the total dry mass content 15.77%, 25.50%, 22.67% and 19.59%, with average mass of about 41 g, 23 g, 36 g and 42g, respectively.

Drying of fruits was carried out in an experimental dryer for testing convective drying process (Kandić et al., 2006). An air streaming drying procedure was used at two constant air temperatures, 90 °C and 70 °C. Control fruits (without pretreatment) and dipped fruits (plum fruits which are, in laboratory conditions, immersed into boiling water for 20 seconds) were dried. Fresh plum fruits were placed in a single layer on an inox tray and 6 trays were placed in a drying chamber. Dipped and control fruits were dried simultaneously in the same experiment on the trays symmetrically placed in the drying chamber (3 trays with the control and 3 trays with dipped fruits). Through the trays with plum fruits, vertical heated air stream with predefined characteristics (temperature, flow) was introduced. Direction of vertical streaming during the process of drying was changed alternately and periodically for 60 minutes, thus achieving the same drying conditions on all of the trays. Fruit drying was completed once the fruits achieve 75% of total dry mater.

Fresh fruits have been kept in plastic bags until chemical analyses at -18 °C, while the dried fruits have been kept in plastic bags at room/ambient temperature (20 °C) until chemical analyses, for conditioning. By testing of fruit chemical composition of the investigated cultivars, the following was determined: content of dry matter (DM), obtained by drying at the laboratory dryer "Sutjeska" (Srbija) at 105 °C until constant mass, and the content of total and invert sugars and sucrose (by Luff-Schoorl's method). Results were expressed in grams per 100 gram of total dry matter (g/100 g DM).

Results were shown as arithmetical mean of three replications \pm standard deviation and were processed statistically using analysis of variance (ANOVA). For testing of the difference significance of mean values of the tested parameters, Duncan's multiple grading test for significance threshold was used $P \leq 0,05$.

Results and Discussion

Table 1 shows the content of total sugars, sucrose, invert sugars in the fresh and dried fruits of the plum cultivars, expressed in g/100 g DM for more accurate and clearer comparison. Percentage of the total sugars, invert sugars and sucrose in dried fruits compared to fresh fruit is shown in Graphs 1, 2 and 3, respectively, with the content of these parameters in fresh fruit shown by a line representing 100% of its content. Percentage of contents of the tested parameters depending on the drying temperature (a) and pretreatment influence (b) are shown separately on Graphs.

Analyzing the contents of total sugars in fresh fruit of the tested cultivars (Table 1), leads to an interesting information that the cultivar 'Čačanska Lepotica' had the highest value of this parameter (68.44 g/100 g DM). This could be explained with the fact that this cultivar, due to the lowest contents of total dry matter (15.77%), compared to the other tested plum cultivars,

has a greatest share of total sugars. In 'Mildora', the content of total sugars was 67.48 g/100 g DM, which was higher in comparison with the cultivars 'Čačanska Rodna' and Stanley, as expected, having in mind that this cultivar is known for the high content of total dry matter (Ogašanović and Ranković, 1996) and the total sugars (Mitrović et al., 2006).

Tab.1. Content of total sugars, invert sugars and sucrose (g/100 g DM) in fresh and dried fruit of cultivars


Cultivar	Sample	Total sugars	Invert sugars	Sucrose
Čačanska Lepotica	Fresh plum	68.44 ± 3.17 a	47.11 ± 2.84	20.26 ± 1.55 a
	Prune			
	90 °C control	59.68 ± 1.56 b	51.61 ± 0.88	7.67 ± 1.56 b
	dipping	57.28 ± 3.84 b	49.66 ± 5.04	7.24 ± 2.35 b
	70 °C control	59.98 ± 3.25 b	54.12 ± 0.10	5.57 ± 3.06 b
dipping	57.60 ± 4.90 b	52.40 ± 6.26	4.93 ± 1.29 b	
Mildora	Fresh plum	67.48 ± 5.08	41.36 ± 3.92 b	24.82 ± 1.58 a
	Prune			
	90 °C control	59.58 ± 2.88	52.07 ± 2.70 a	7.13 ± 0.95 b
	dipping	60.25 ± 4.45	51.21 ± 3.01 a	8.59 ± 1.61 b
	70 °C control	62.31 ± 3.26	55.31 ± 1.54 a	6.62 ± 1.74 b
dipping	60.56 ± 3.05	54.67 ± 2.34 a	5.60 ± 0.88 b	
Čačanska Rodna	Fresh plum	65.57 ± 1.66 a	39.27 ± 2.76 b	24.97 ± 3.71 a
	Prune			
	90 °C control	55.84 ± 2.59 b	47.52 ± 4.41 a	7.90 ± 1.92 b
	dipping	55.33 ± 2.95 b	48.07 ± 2.88 a	6.89 ± 1.63 b
	70 °C control	58.23 ± 2.47 b	47.54 ± 2.81 a	10.15 ± 2.87 b
dipping	58.67 ± 3.51 b	50.19 ± 2.47 a	8.06 ± 2.78 b	
Stanley	Fresh plum	64.80 ± 0.99 a	39.67 ± 1.77 b	23.86 ± 2.20 a
	Prune			
	90 °C control	55.71 ± 3.29 b	48.53 ± 1.74 a	6.83 ± 1.67 b
	dipping	54.18 ± 2.43 b	48.78 ± 1.94 a	5.13 ± 0.48 b
	70 °C control	57.00 ± 1.26 b	48.35 ± 1.55 a	8.22 ± 0.49 b
dipping	57.20 ± 1.33 b	48.50 ± 1.71 a	8.26 ± 1.56 b	

Data followed by different letters within each column and cultivar are significantly different according to Duncan's multiple range test at $P \leq 0.05$

In dried fruits of all tested cultivars, except in 'Mildora', a lower content of total sugars was found, compared to initial fresh fruit regardless of the applied drying temperature and dipping procedure (Table 1, Graph 1). That loss is about 10-15%, although there was no juice leakage during the drying process. This was probably because of the participation of sugars in non-enzymatic browning reactions. Wilford et al. (1997), investigating the kinetics of sugar change during the drying of the plum cultivar d'Agen at three drying temperatures using convective drying, concluded that the Maillard reaction starts at about 45-50% of moisture loss, and the proof for that is profile change in the presence of some sugars.

During the drying process, there comes to the hydrolysis of sucrose in plum fruit and the increase in the contents of invert sugars. Analyzing Graph 2, dried fruit of all tested cultivars were found to act similarly regardless of the drying temperatures and dipping procedure applied, i.e. the increase is about 25-30%. Exception is the plum cultivar 'Čačanska Lepotica', in which a minimum increase in the content of invert sugars was observed, of no statistical


importance, and which was lower in fruits dried at higher temperatures (7.5% in total) than in dipped fruits where the increase is only 8.3%.



Graph. 1. Total sugars content in prunes compared to the fresh plums, expressed as percentages

Legend:

- Cultivar: 1- Čačanska Lepotica; 2- Mildora; 3- Čačanska Rodna; 4- Stanley
- Influence of the drying temperature: 90 °C; 70 °C
- Influence of the dipping pretreatment: K- control; D - dipping



Graph. 2. Invert sugars content in prunes compared to the fresh plums, expressed as percentages

Legend:


- Cultivar: 1- Čačanska Lepotica; 2- Mildora; 3- Čačanska Rodna; 4- Stanley
- Influence of the drying temperature: 90 °C; 70 °C
- Influence of the dipping pretreatment: K- control; D - dipping

Examining the carbohydrate content of prunes and their products, Dikeman et al. (2004) reported a significant increase in the content of glucose and fructose, the two most important monosaccharides in plums, in relation to fresh fruit as a starting material. Cinquanta et al.

(2002) found the increased content of glucose and fructose in dried fruits of all the plum cultivars, whereas in the cultivar 'Stanley' differences in the increased content of these simple sugars among fruits dipped in different manners were noted to be of no statistical significance, compared with non-treated fruits.

As formerly stated, results of the sucrose content in fresh and dried fruit of the investigated plum cultivars are shown in Table 1. In all tested cultivars, dramatically lower sucrose content in dried plum was found compared to fresh fruit.

Analyzing Graph 3, drying temperature is found to have greater impact on the reduced sucrose content in dried fruit of all tested cultivars in comparison with the applied pretreatments. Dried fruits of the cultivars 'Čačanska Lepotica' and 'Mildora' acted in similar way, sucrose content in fruits dried at 90 °C was about 3 times lower than in fresh fruits, while in fruits dried at 70 °C, the content was 4 times lower. In cultivars, 'Čačanska Rodna' and 'Stanley' the opposite was found, and a lower sucrose loss was observed by a lower drying temperature.



Graph. 3. Sucrose content in prunes compared to the fresh plums, expressed as percentages

Legend:

- Cultivar: 1- Čačanska Lepotica; 2- Mildora; 3- Čačanska Rodna; 4- Stanley
- Influence of the drying temperature: 90 °C; 70 °C
- Influence of the dipping pretreatment: K- control; D - dipping

Dipping as an applied pretreatment, had much less effect on the reduced sucrose content in dried fruit compared with non-treated fruits, i.e. control. A slight difference was observed between dipped and control fruits in decreased sucrose content in the cultivars 'Čačanska Lepotica' and 'Stanley', whereas in 'Mildora', which dried fruits contain about 3.5 times less sucrose compared to the starting raw material, sucrose loss is the same regardless of whether the fruits are dipped or not. Only in the plum cultivar 'Čačanska Rodna', effect of dipping on reduced sucrose content was recorded so that the dipped fruits contain 3.3 times less sucrose than the fresh fruits and in non-treated fruits, representing control, sucrose amount was reduced 2.7 times.

During the process of plum drying, the hydrolysis of sucrose occurs, thereby generating glucose and fructose. Analyzing the kinetics of carbon hydrate change during drying of the plum cultivar d'Agén, Wilford et al. (1997) found that, during the drying process, sucrose is completely hydrolysed. According to them, low pH value and high moisture of fruit in early stages of drying favour the hydrolysis of sucrose generating glucose and fructose. In their

investigations, at the drying temperature of 70 °C, sucrose was completely vanished in 6-7 hours, while at 90 °C, sucrose loss was noted after 2 hours of drying. Authors Di Matteo et al. (2003) also noted a complete loss of sucrose in dried fruit of the plum cultivar Angeleno. Cinquanta et al. (2002) recorded the complete sucrose loss in dried fruit of the cultivars Angeleno and Empress too, while on the other side in the cultivar 'Stanley' only reduced content of sucrose in dried fruit compared with fresh fruit was noted, thus emphasizing that 'Stanley' is an excellent cultivar for drying. Decrease in sucrose content in the course of drying is not a plum characteristic only as a fruit species. Moreover, in the course of banana drying, the hydrolysis of sucrose occurs, i.e. there is a decrease in content of sucrose in comparison with the starting raw material. (Leite et al., 2007).

Conclusion

Based on the investigation results of the drying temperature and dipping effect on the change of total sugars, invert sugars and sucrose content in fruit of the plum cultivars 'Čačanska Lepotica', 'Mildora', 'Čačanska Rodna' and 'Stanley', the following conclusions can be drawn:

- Drying temperature affected the change of total sugars, invert sugars and sucrose content in fruit of the investigated plum cultivars. Change intensity was conditioned by varietal characteristics.
- Dipping, as a pretreatment applied, had no effect on the change of these parameters content, except in the cultivar 'Čačanska Rodna' in which greater decrease of sucrose compared to non-treated fruits was noted.
- In dried plum fruits of the cultivars 'Čačanska Lepotica', 'Čačanska Rodna' and 'Stanley', a lower content of total sugars was noted, based on the calculation of the total dry matter in relation to initial fresh fruit, regardless of the drying temperatures and dipping procedure applied. The only exception was the cultivar 'Mildora' in which there were no statistical differences in the total sugar content in dried fruit compared to fresh fruit.
- During the drying process, an increase of invert sugars content occurred in all tested cultivars, except in 'Čačanska Lepotica', in which there was no statistically significant increase in the content.
- In the course of plum drying, the hydrolysis of sucrose occurred, which was manifested in a dramatic decrease of its content in dried fruits compared to starting raw material in all tested cultivars.

Acknowledgements

The study was financed from the funds of the Ministry of Education, Science and Technological Development of the Republic of Serbia (project TR-31093 and OI-172057).

References

- Cinquanta L., Di Matteo M., Esti M. (2002): Physical pretreatment of plums (*Prunus domestica*). Part 2. Effect on the quality characteristics of different prune cultivars. *Food Chemistry*, 79: 233-238.
- Di Matteo M., Cinquanta L., Galiero G., Crescitelli S. (2003): A mathematical model of mass transfer in spherical geometry: plum (*Prunus domestica*) drying. *Journal of Food Engineering*, 58: 183-192.
- Dikeman C. L., Bauer L. L., Fahey G. C. (2004): Carbohydrate Composition of Selected Plum/Prune Preparations. *Journal of Agricultural and Food Chemistry*, 52, 4: 853-859.
- Friedman M. (1996): Food Browning and Its Prevention: An Overview. *Journal of Agricultural and Food Chemistry*, 44, 3, 631-649.
- Kandić M., Mitrović O., Gavrilović -Damnjanović J. and Mitrović V. (2006). Flow-through model for the assessment of fruit drying procedure. *Voćarstvo* 40, 379-388.
- Kim J.S., Lee Y.S. (2008): Effect of reaction pH on enolization and racemization reactions of glucose and fructose on heating with amino acid enantiomers and formation of melanoidins as result of the Maillard reaction. *Food Chemistry*, 108: 582-592.
- Leite J. B., Mancini M. C., Borges S. V. (2007): Effect of drying temperature on the quality of dried bananas cv. *Prata* and *d' àqua*. *LWT*, 40: 319-323.
- Manzocco L., Calligaris S., Mastrocola D., Nicoli M. C., Lerici C. R. (2001): Review of non-enzymatic browning and antioxidant capacity in processed foods. *Trends in Food Science and Technology*, 11: 340-346.
- Mitrović O., Gavrilović-Damnjanović J., Popović B., Kandić M. (2006): Properties of Čačak plum cultivars suitable for drying. *Voćarstvo*, 40, 155: 255-261.
- Ogašanović D., Ranković M. (1996): Major characteristics of some plum hybrids resistant to sharka. *Journal of Yugoslav Pomology*, 30, 113/114: 117-122. (in Serbian)
- Piga A., Del Caro A., Corda G. (2003): From plums to prunes: Influence of drying parameters on polyphenols and antioxidant activity. *Journal of Agricultural and Food Chemistry*, 51: 3675-3681.
- Popović B., Nikićević N., Gavrilović-Damnjanović J., Mitrović O., Petrović A., Ogašanović D. (2006): Technological properties of fruits of more important plum cultivars used as a raw material for plum brandy production. *Arhiv za poljoprivredne nauke*, 67, 238: 73-82.
- Sanz M.L., del Castillo M.D., Corzo N., Olano A. (2001): Formation of amadori compounds in dehydrated fruits. *Journal of Agricultural and Food Chemistry*, 49: 5228-5231.
- Zlatković B. (2000): The role of processing technology on the plum marketing. *Tematic Proceedings of 1st International Scientific Symposium: Production, Processing and Marketing of Plums and Plum products*, Koštunići (Republic of Serbia), 245-252. (in Serbian)
- Wilford L. G., Sabarez H., Price W.E. (1997): Kinetics of carbohydrate change during dehydration of d'Agen prunes. *Food Chemistry*, 59, 1: 149-155.